

# THE STAR FORMATION HISTORY OF THE LOCAL GROUP

Eva K. Grebel<sup>1,2,3</sup>

<sup>1</sup>University of Washington, Astronomy Department, Box 351580, Seattle, WA 98195, USA

<sup>2</sup>Hubble Fellow

<sup>3</sup>Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

## ABSTRACT

The star formation histories of Local Group galaxies are summarized. The three large spirals are discussed individually. The discussion of the Local Group dwarfs concentrates on differences and commonalities. While Local Group galaxies exhibit an amazing diversity in their star formation and enrichment histories even within the same morphological type, they share a number of common global properties. Major determining factors for their evolution are galaxy mass and environmental effects such as ram pressure and tidal stripping. It seems likely that low-mass dwarf irregular galaxies evolve into dwarf spheroidals.

Key words: Local Group – dwarf galaxies – galaxy evolution – star formation – interstellar medium

## 1. INTRODUCTION

Our Milky Way and its companions are part of a sparse group of galaxies known as the Local Group (LG). Due to the uncertainties in the mass estimates for LG galaxies and the lack of information about galaxy orbits the LG boundaries are poorly defined. The LG zero-velocity surface has a radius of  $\sim 1.2$  Mpc (Courteau & van den Bergh 1999) around the LG barycenter when a simplified spherical potential is adopted. Within this zero-velocity surface there are currently 36 galaxies known, most or all of which may be members of the LG. This radius definition excludes galaxies such as SagDIG or the nearby NGC 3109/Sextans minigroup (van den Bergh 1999a), which has a mean barycentric distance of 1.6 Mpc from the LG and is kinematically offset from the  $\pm 60$  km s<sup>-1</sup> velocity dispersion envelope of the LG. In contrast to earlier definitions (Grebel 1997; Mateo 1998) this radius also avoids overlap with the closest members of the elongated Sculptor group, which are located at  $\gtrsim 1.7$  Mpc. When viewed on a larger scale, the Local Group appears to be one of several condensations in a clumpy filament that extends over many megaparsecs and contains a number of nearby groups and clusters (Tully & Fisher 1987).

## 2. GALAXY TYPES IN THE LOCAL GROUP

The absolute magnitudes of LG galaxies range from  $\sim -21 < M_V < -8.5$  mag and comprise masses of a few  $10^{12} M_\odot$  to  $10^7 M_\odot$ . As in any galaxy group and cluster the majority of the galaxies in the LG are dwarf galaxies. If we consider all galaxies with  $M_V > -18$  mag to be dwarf galaxies, then the LG contains 31 dwarfs. Three of the four galaxies that are more luminous than this cutoff are spirals (the Milky Way, M31, M33), and one is a gas-rich, active irregular galaxy (the Large Magellanic Cloud (LMC)). The LMC is the closest irregular satellite of the Milky Way, and M33 is part of the M31 subgroup.

The LG dwarfs can be subdivided in three basic types: dwarf irregulars (dIrrs), dwarf ellipticals (dEs), and dwarf spheroidals (dSphs). Note that these morphological distinctions are arbitrary to some extent and are used in differing ways by different authors. In the following the term “old population” refers to ages  $> 10$  Gyr, “intermediate age” denotes ages from 1 Gyr to 10 Gyr, and “young” includes all stellar populations below 1 Gyr.

### 2.1. LOCAL GROUP DWARF IRREGULARS

dIrrs tend to be gas-rich, irregularly shaped galaxies that show scattered, ongoing star formation while spiral density waves are absent. Most dIrrs contain stellar populations of all ages as well as neutral and molecular gas and dust. Some of the LG irregulars/dIrrs show evidence for a bar (e.g., LMC, NGC 6822), while in the Small Magellanic Cloud (SMC) the apparent presence of a stellar bar may be feigned by highly irregular recent star formation (Zaritsky et al. 2000).

### 2.2. LOCAL GROUP DWARF ELLIPTICALS

dEs stand out through their distinctive elliptical shape and compact, dense cores. Their dominant stellar populations are of old and intermediate age, but recent star formation is observed as well. As pointed out by van den Bergh (1999b) and references therein, the structural parameters of the M31 companion M32 make this dwarf galaxy a genuine elliptical rather than a dE/dSph. The remaining three dEs (NGC 205, NGC 185, and NGC 147; all M31 satellites) follow similar scaling relations as the LG dSphs (e.g., Ferguson & Binggeli 1994) and are not

Table 1. A summary of global properties shared by many of the three main dwarf galaxy types in the Local Group. “R” denotes typical tidal radii. “SFR” refers to the present-day star formation rate. Individual dwarf galaxies may differ from the listed values. For comparison the inferred properties of compact high velocity clouds (CHVCs) are also listed.

| Type  | $M_V$<br>[mag] | $\mu_V$<br>[mag arcsec $^{-2}$ ] | R<br>[kpc]    | SFR<br>[ $M_\odot \text{ yr}^{-1}$ ] | $M_{\text{HI}}$<br>[ $M_\odot$ ] | $M_{\text{tot}}$<br>[ $M_\odot$ ] |
|-------|----------------|----------------------------------|---------------|--------------------------------------|----------------------------------|-----------------------------------|
| dIrr  | $\gtrsim -18$  | $\lesssim 23$                    | $\lesssim 5$  | $\lesssim 0.01$                      | $\lesssim 10^9$                  | $\lesssim 10^{10}$                |
| dE    | $\gtrsim -17$  | $\lesssim 21$                    | $\lesssim 4$  | 0                                    | $\lesssim 10^8$                  | $\lesssim 10^9$                   |
| dSph  | $\gtrsim -14$  | $\gtrsim 22$                     | $\lesssim 3$  | 0                                    | $\lesssim 10^5$                  | $\sim 10^7$                       |
| CHVCs | ?              | ?                                | $\lesssim 10$ | 0?                                   | $\lesssim 10^7$                  | $\gtrsim 10^8$                    |

always distinguished from dSphs in the literature. Little to no gas (H I and CO) has been detected in LG dEs (e.g., Sage et al. 1998). Both M32 and NGC 205 have a luminous semi-stellar core and may be considered nucleated (dE,N). M32 appears to contain a massive central black hole (van der Marel et al. 1997).

### 2.3. LOCAL GROUP DWARF SPHEROIDALS

DSphs as defined by Gallagher & Wyse (1994) do not show nuclei and have very little central concentration. They are the least massive and least luminous galaxies known. Their stellar populations are predominantly of old and intermediate age. The lack of H I at the optical positions of these galaxies not understood. DSphs, once thought to be an extension of globular clusters, differ from globulars in several crucial ways: (1) they have very low stellar densities even in their centers, (2) most of them have experienced multiple or extended episodes of star formation, (3) they have high mass-to-light ratios (Mateo 1998, but see Klessen & Kroupa 1998 for a different view), and (4) some dSphs contain globular clusters themselves. Except for Tucana and Cetus, LG dSphs are close companions of either the Milky Way or M31 (Fig. 1).

A summary of representative dwarf properties in the LG is given in Table 1. For further discussion of morphological types in and around the Local Group we refer to Grebel (1999) and the very detailed reviews by van den Bergh (1999b, 2000a). Globular clusters have been identified in all galaxy types in the LG, though only 13 LG galaxies contain globulars (see Grebel 2000 for a review).

### 2.4. COMPLETENESS AND MISSING SATELLITES

Many of the faint LG members were discovered only in recent years. It is conceivable that the galaxy census continues to be incomplete for faint dwarf galaxies and in regions of high foreground extinction. Destruction of dwarfs through accretion by spirals may have played an important role in the past and is currently evidenced by the merger of the Sagittarius dSph galaxy with the Milky Way (Ibata et al. 1994). A comparison with the luminos-

ity functions of nearby groups (Trentham 1998) suggests that the LG census is still deficient. Deep wide-field surveys such as the Digitized Second Palomar Sky Survey (Djorgovski et al. 1999) and the Sloan Digital Sky Survey (Gunn & Weinberg 1995) may lead to the detection of additional LG dwarfs of very low surface brightness.

Hierarchical cold dark matter (CDM) models predict roughly a factor 10 more dark matter halos for the LG than the number of known dwarf galaxies (Klypin et al. 1999; Moore et al. 1999). If the CDM model predictions are correct and if these dark halos have baryonic counterparts, then compact high velocity clouds (CHVCs; Braun & Burton 1999) may be good candidates for the predicted LG satellites. While most high-velocity cloud complexes are within 10 kpc of the Milky Way or part of the Magellanic stream, CHVCs appears to be at distances of up to 1 Mpc, show infall motion with respect to the LG barycenter, have masses of  $\sim 10^7 M_\odot$ , are rotationally supported, and dark matter dominated (Braun & Burton 2000). So far it is unknown whether CHVCs have stellar counterparts or might constitute low-mass protogalaxies prior to star formation.

### 2.5. MORPHOLOGICAL SEGREGATION

The galaxy distribution in the LG is dominated by two prominent subgroups around its two most massive members, the Milky Way and M31 (Fig. 1). 70% of the smaller galaxies are found within  $\sim 300$  kpc around the two dominant Sb/Sbc spirals. Wilkinson & Evans (1999) suggest that more than 50% of the LG mass is concentrated around the two large spirals. There are no massive ellipticals in the LG, and the third spiral, M33, does not have any known companions. The dependence of distribution within the LG on galaxy type is also known as morphological segregation and may hold clues to the impact of environmental effects on the evolution of dwarf galaxies in groups. Morphological segregation is also evident in the nearby M81 group (Karachentsev et al. 2000). On a much larger scale similar effects may be seen in the morphology-density re-




Figure 1. Number distribution of Local Group galaxies as a function of distance. Morphological segregation is evident. Upper panel: Dwarf ellipticals and dwarf spheroidal galaxies (the least massive dwarfs) are mostly found in close proximity of M31 or the Milky Way. Middle panel: Dwarf irregular galaxies are more uniformly distributed throughout the Local Group. The less massive galaxies among the dwarf irregulars with  $250 < D_{\text{Spiral}} < 450$  kpc are often referred to as transition-type objects (see text). Lower panel: Local Group galaxies tend to cluster around the two massive spirals. Local Group membership for the outlying objects at larger distances is uncertain.

lation in galaxy clusters at low and high redshift (Dressler 1980; Dressler et al. 1997).

### 3. DERIVING STAR FORMATION HISTORIES

#### 3.1. STATE OF THE ART: SYNTHETIC CMD TECHNIQUES

The derivation of star formation histories of resolved stellar populations relies heavily on photometric imaging and color-magnitude diagrams (CMDs). The most detailed star formation histories are determined by modelling observed CMDs through the best-matched synthetic CMD (see Aparicio 1999 for a review of synthetic CMD techniques). The results depend on the quality and depth of the photometric input data, the treatment of observational errors, the input evolutionary models, the sophistication of the modelling codes, etc. Free parameters include the choice of an IMF, binary fraction, and enrichment law. Due to different approaches and differing parameter choices results obtained by different groups may not be directly compa-

table. Some of the problems in deriving reliable star formation histories both on the theoretical and observational side are outlined below.

#### 3.2. SHORTCOMINGS OF EVOLUTIONARY MODELS

While very impressive progress has been made, models are limited by how well we understand the details of stellar evolution (e.g., mass loss in different phases of stellar evolution, the impact of rotation and mixing, life times, blue-to-red supergiant ratios, the second-parameter problem, etc.). The currently used models are “unphysical” in the sense that they were calculated for non-rotating, single stars. When the new rotational models become available it will still be difficult to disentangle effects of rotation and binarity in observational data. Problems also persist in transforming theoretical models to the observational plane, which can lead to discrepancies especially for cool stars (see Fig. 5 in Grebel 1999).

#### 3.3. LACK OF ABUNDANCE INFORMATION

Broadband data (usually obtained in only two filters) are insufficient to break the age-metallicity degeneracy particularly when dealing with complex, mixed populations. Metallicity-sensitive photometric systems work fairly well in populations with little age spread and can efficiently provide abundances (though not necessarily iron abundances) for a large number of stars. Spectroscopy is considered the most reliable method for abundance determinations, but it is disquieting to consider the wide variety of spectroscopic abundances found by various authors for the extensively studied young SMC cluster NGC 330 (see González & Wallerstein 1999).

For most LG galaxies spectroscopic abundance information has either not been obtained or is available only for very few evolutionary phases.  $[\text{Fe}/\text{H}]$  estimates are typically from the infrared Ca II triplet of old red giants, from integrated spectra of globular clusters, or from luminous supergiants. Photometric abundances are usually based on the photometric colors and slope of the red giant branch (Armandroff & Da Costa 1990). Oxygen abundance measurements exist for planetary nebulae (PNe) or H II regions in about half of the LG galaxies (Mateo 1998). The relation between nebular abundances from H II regions and PNe and stellar  $[\text{Fe}/\text{H}]$  values is poorly understood (Richer & McCall 1995). Discrepancies remain even when comparing element abundances of young objects such as massive stars and H II regions (e.g., Garnett 1999).

#### 3.4. AREA COVERAGE

Most LG galaxies show evidence for spatial variations in their star formation histories. The sampling of small subsections of a galaxy may not be representative for the overall star formation history or lack the number statistics to

detect less well-populated age ranges (see Grebel 1997). This affects in particular old populations. In more distant dIrrs old populations were detected through deep ground-based observations of their less crowded halos (Minniti & Zijlstra 1996; Minniti et al. 1999). Deep *HST* data of the center of Leo I led to the suggestion of a lacking old population in this dSph (Gallart et al. 1999), while wide-area imaging revealed a well-populated old horizontal branch (Held et al. 2000).




Figure 2. With increasing distance it becomes difficult to reach the main-sequence turnoff of the older populations, which is one of the factors that contribute to decreased resolution for older ages. In dIrrs superimposed younger main sequence stars aggravate the problem and may not only make the oldest main-sequence turnoff and subgiant branch hard to measure but also merge with the old blue horizontal branch.

### 3.5. AGE RESOLUTION

While we can determine the ages for young populations traced by massive stars to a few  $10^6$  to  $10^7$  years, the age resolution decreases roughly logarithmically for older populations (see Fig. 4 in Grebel 1999) within a given galaxy. Furthermore, the attainable age resolution decreases with distance from the Milky Way (Fig. 2).

## 4. STAR FORMATION IN THE THREE SPIRALS

A detailed description of the star formation histories of the three spirals in the LG is beyond the scope of this

short paper. Instead I will attempt a brief summary and refer to the recent reviews by Majewski (1999) and van den Bergh (1999b; 2000a) for more information.

The star formation histories of spiral galaxies vary with position within the galaxy depending on which galaxy component is considered – halo, bulge, or disk, and further morphological or kinematic subdivisions.

### 4.1. THE MILKY WAY

The Milky Way is a S(B)bc galaxy with continuing star formation in the Galactic center (Figer et al. 1999) and in its spiral arms. The oldest components of the Milky Way are the old halo (both field stars and old globulars) and the thick disk. The formation of the old halo may be best described by Searle & Zinn (1978)-like accretion. Most of the merger events are believed to be ancient (Unavane et al. 1996). Efforts are on the way to trace their remnants by surveying the Galactic halo for kinematic substructure (e.g., Morrison et al. 2000).

The earliest significant star formation in the thick disk may date back 13 Gyr ago. The thick disk has a mean metallicity of  $-0.6$  dex (Majewski 1999) and a dynamically hot, extended structure. The thin disk experienced multiple bursty star formation episodes starting  $\sim 9$  Gyr ago. Rocha-Pinto et al. (2000) conclude that the global disk star formation rate in the Milky Way fluctuates on time scales of less than  $0.2 - 1$  Gyr with amplitudes exceeding a factor  $2 - 3$ . They suggest that such intermittent star formation may be typical for disk galaxies though for the Milky Way interactions with the Magellanic Clouds may have played a role as well. While there are indications of a radial abundance gradient in the thin disk the metallicity is a stronger function of galactocentric radius than of age and exhibits large scatter at any position and age (Edvardsson et al. 1993). Mean metallicities range from  $-0.5 < [\text{Fe}/\text{H}] < 0.1$  dex with a pronounced metal-poor tail. One of the scenarios for thin-disk evolution is inside-out formation (Edvardsson et al. 1993), i.e., the inner parts form more rapidly and experience stronger enrichment.

The Galactic bulge as well as other bulges seem to have undergone early and very rapid enrichment. The mean metallicity of the bulge is  $-0.25$  dex with a spread that can easily exceed 1 dex (Minniti et al. 1995). Also, a radial abundance gradient is observed. Both the abundance spread and wide range in ages may be in part caused by contributions from overlapping Milky Way halo and disk components. Bulge formation in a rapid monolithic collapse similar to the scenario suggested by Eggen et al. (1962) appears likely.

### 4.2. M31

M31, an Sb I–II spiral, usually believed to be  $\sim 30\%$  more massive than the Milky Way (Wilkinson & Evans 1999),

may in fact be only as massive or even less massive than the Milky Way (Evans et al. 2000). It shows on average higher metallicities than the Milky Way and a considerable abundance spread. The outer halo has a mean metallicity of  $-1.2$  dex, while for the inner halo  $\langle[\text{Fe}/\text{H}]\rangle = -1.0$  dex (van den Bergh 2000a and references therein). M31 appears to have undergone rapid enrichment as a whole. The bulge emits  $\sim 30\%$  of the visible light of M31 and contains two nuclei, which may be both part of the same eccentric disk of stars orbiting a central black hole (Kormendy & Bender 1999). M31 has a warped stellar and gas disk, possibly caused by tidal interactions with massive nearby companions such as M32. The horizontal branch morphology of some of the  $\sim 600$  globular clusters in M31 suggests that they have similarly ages as the oldest Galactic globulars (e.g., Ajhar et al. 1996). M31 currently only shows low star formation activity. The dominant present-day star formation is occurring in the outer disk. The UV line strengths of massive OB stars in M31 indicate a metallicity comparable to that of the young Milky Way population (Bianchi et al. 1996).

#### 4.3. M33

M33 is an Sc II–III spiral with a significant radial abundance gradient. It shows a nucleus, a pronounced disk and halo, but no obvious bulge. The mean metallicity of M33's halo ( $-1.6$  dex) is lower than M31's and shows a spread. Stellar and H I disk are tilted with respect to each other, but M33 doesn't have any known companions that might be responsible. About 54 globular clusters are currently known in M33 (Mochejska et al. 1998). The lack of a blue horizontal branch (BHB) in M33 globulars may suggest that the bulk of M33's star formation began several Gyr later than in the Milky Way (Sarajedini et al. 1998). In the past  $10^7$  to  $10^8$  years cluster formation increased (Chandar et al. 1999) and appears to be correlated with gas inflow into the center of M33. The UV line strengths of massive OB stars in M33 resemble the young LMC population in metallicity (Bianchi et al. 1996).

### 5. STELLAR LUMINOSITY FUNCTIONS

The behavior of the high-mass portion of the stellar luminosity function (LF) and initial mass function (IMF) is directly measurable only in well-resolved young populations. The careful spectroscopic and photometric studies by Massey et al. (1995a,b) of OB associations in the Milky Way, LMC, and SMC showed that the upper IMFs vary little and are essentially consistent with a Salpeter IMF within the errors despite the differences in metallicity. This was confirmed for a range of masses down to  $2.8 - 0.7 M_{\odot}$  for young clusters in the LMC (Massey & Hunter 1998; Elson et al. 1999; Grebel & Chu 2000). The only known exceptions in single-age populations are the Galactic center clusters and the central starburst cluster in the

Galactic giant H II region NGC 3603, which seem to have significantly flatter IMFs (Figer et al. 1999; Eisenhauer et al. 1998).

The LFs of older, low-mass field populations in the Galactic bulge (Holtzman et al. 1998), LMC and SMC (Holtzman et al. 1999b), Draco (Grillmair et al. 1998), and Ursa Minor (Feltzing et al. 1999) are in excellent agreement with the solar neighborhood IMF or that of globular clusters unaffected by significant mass segregation.

Apart from providing support for the universality of the IMF the findings for dSphs show that low-mass stars down to masses of  $0.45 M_{\odot}$  are not responsible for the inferred high mass-to-light ratios of these galaxies.

### 6. THE OLDEST POPULATIONS IN THE LOCAL GROUP

Old populations ( $> 10$  Gyr) have been detected in all LG galaxies studied in sufficient detail. Tracers of the oldest populations include old main-sequence turnoffs and subgiant branches, horizontal branches, and RR Lyrae in globular clusters and field populations.

Deep imaging largely based on *HST* data has established that a number of LG galaxies share a common epoch of ancient star formation. Main-sequence photometry reveals that the oldest globular clusters in the Milky Way halo and bulge, in the LMC (Olsen et al. 1998), in Sagittarius (Montegriffo et al. 1998), and in Fornax (Buonanno et al. 1998) are coeval within the measurement accuracy. Similarly, deep main-sequence photometry of the oldest field populations in Sagittarius (Layden & Sarajedini 2000), LMC (Holtzman et al. 1999a), Draco, Ursa Minor (Mighell & Burke 1999), Fornax, Sculptor (Monkiewicz et al. 1999), Carina (Mighell 1997), and Leo II (Mighell & Rich 1996) indicates that they have the same relative age as the oldest Galactic globular clusters.

Though main-sequence photometry is lacking, the CMD of the globular cluster in the WLM dIrr (Hodge et al. 1999), the BHBs of globulars in M31 (Ajhar et al. 1996) and in NGC 147 (Han et al. 1997), and spectroscopy of one of NGC 6822's globulars (Cohen & Blakeslee 1998) are interpreted as indicative of ages similar to those of the old Galactic globular clusters. Comparatively old ages may be implied by the existence of a BHB in the field populations of Sextans (Mateo et al. 1991), Leo I (Held et al. 2000), Phoenix (Smith et al. 2000), IC 1613 (Cole et al. 1999), Cetus (Tolstoy et al. 2000), And I (Da Costa et al. 1996), And II (Da Costa et al. 2000), NGC 185 (Geisler et al. 1999), NGC 147 (Han et al. 1997), and Tucana (Lavery et al. 1996). However, while a BHB indicates an old population it should be emphasized that due to second-parameter effects other than age a BHB does not necessarily imply an ancient, old halo globular-cluster-like population.

Conversely (and with the same caveat), the absence of a BHB in M33 globulars (Sarajedini et al. 1998) and the apparent lack of a pronounced BHB in the field populations of the SMC, WLM (Dolphin 2000), Leo A (Tolstoy

et al. 1998), and DDO 210 (Tolstoy et al. 2000) may indicate that the bulk of the old population in these galaxies formed with a delay of a few Gyr.

## 7. SPATIAL VARIATIONS AND TIDAL EFFECTS

### 7.1. SPATIAL VARIATIONS IN STAR FORMATION HISTORY

Spatial variations in star formation history are a common occurrence in galaxies (see also Section 3.4) and provide clues about the distribution of the star-forming material as a function of time, the dynamical history of galaxies as well as of the gas retention or gas loss history. The extent of spatial variations seems to be correlated both with the overall mass of the galaxy and its environment.

A global property of LG galaxies is that the oldest populations are the most extended ones. This may indicate that galaxies have contracted with time (van den Bergh 1999b) or may be a consequence of dispersion or substantial mass loss. High-mass dwarf galaxies such as dIrrs typically show multiple distinct zones of concurrent star formation. Some of these regions can be long-lived, exist for several 100 Myr, and may migrate (e.g., Sextans A: Dohm-Palmer et al. 1997; LMC: Grebel & Brandner 1999). In contrast, low-mass satellites (especially dSphs) tend to have centrally concentrated intermediate-age or young populations and may show radial age gradients. Star-forming material appears to have been retained longer in the central regions. An example of a dSph with a very pronounced age and metallicity gradient is Fornax (Stetson et al. 1998; Grebel & Stetson 1999). In Fornax different populations have slightly different centroids, and the youngest population is asymmetrically distributed.

In some cases spatial variations in star formation history are traced only by horizontal-branch variations in the sense that BHB stars are more spatially extended than red horizontal branch stars (examples include Sculptor (Hurley-Keller et al. 1999) and And I (Da Costa et al. 1996)). A few low-mass dwarf galaxies show no apparent indication for spatial population variations including Ursa Minor, Carina (Smecker-Hane et al. 1996), and And II (Da Costa et al. 2000).

### 7.2. SPATIAL VARIATIONS IN STELLAR DENSITY

While lacking variations in its stellar populations, Ursa Minor shows statistically significant stellar density variations (Kleyna et al. 1998) instead of a smooth surface density profile. In the more luminous dSph Fornax four out of the five globular clusters are found at distances larger than the galaxy's core radius although their orbital decay scales through dynamical friction are only a few Gyr (Oh et al. 2000), a fraction of the globular clusters' ages. Significant mass loss through Galactic tidal perturbation and the resulting decrease in the satellite galaxy's gravitational potential may have increased the clusters' orbital

semimajor axes and efficiently counteracted the spiralling-in through dynamical friction (Oh et al. 2000).

The currently available radial velocity measurements for individual stars in dSphs do not yet allow one to distinguish between dynamical models where mass follows light, extended dark halo models, and disrupted remnant models without dark matter (Kleyna et al. 1999). Interestingly, the detection of a possible very extended population of extratidal stars around the dSph Carina might imply that this galaxy has now been reduced to a mere 1% of its initial mass (Majewski et al. 2000), the rest of which has already been accreted by the Milky Way's halo. Apart from favoring disrupted remnant models this claim has far-reaching consequences for the determination of star formation histories of nearby low-mass satellites: If they are indeed significantly tidally disrupted then their present-day stellar content cannot easily be used to derive evolutionary histories over a Hubble time. The observed metallicity spread in the nearby dSph Draco and other dSphs provides further evidence for significant past mass loss since this dwarf was once massive enough to retain its metals for subsequent star formation. According to Mac Low & Ferrara (1999) almost no metals are retained by galaxies with  $< 10^9 M_{\odot}$ .

Substantial tidal disruption by the Milky Way is evidenced by the Magellanic Clouds and Magellanic Stream as well as by the Sagittarius dSph. Tracer features include gaseous and stellar tidal tails, kinematics, and structural parameters (e.g., Hatzidimitriou et al. 1993, Putman et al. 1998, Majewski et al. 1999, Weinberg 2000, Ibata et al. 2000). The dSphs Draco and Ursa Minor as well as various globular clusters may be part of the extended Magellanic Stream (Kunkel 1979). Peaks in the star cluster formation rate could be related to close encounters between the Magellanic Clouds and the Milky Way (Girardi et al. 1995, Grebel et al. 1999). The stellar metallicity patterns in the intercloud region impose time constraints on the tidal disruption (Rolleston et al. 1999).

### 7.3. VARIATIONS IN THE GAS CONTENT OF DWARFS

The gas in dwarf galaxies shows density and spatial variations as well.

The H I in dIrrs tends to be even more extended than the oldest stellar populations and shows a clumpy distribution. Central H I holes or off-centered gas concentrations are observed in a number of cases and may be driven by star formation events (Young & Lo 1996; 1997a,b). Ongoing gas accretion has been found in at least one dIrr (IC 10: Wilcots & Miller 1998). Low amounts of gas ( $\sim 10^5 M_{\odot}$ ) were detected in dEs (Sage et al. 1998). The gas kinematics of NGC 185 indicate an internal origin (i.e., consistent with stellar gas loss), while the significant angular momentum of the H I in NGC 205 as compared to the non-rotating stellar component may imply accretion. Single-dish observations of dSphs did not detect gas (e.g.,



Knapp et al. 1978), and subsequent VLA synthesis studies covering much of the optical extent of some of the dSphs did not reveal HI in emission or in absorption down to column densities of  $10^{17} - 10^{18} \text{ cm}^{-2}$  (Young 1999; 2000).

The apparent absence of gas in dSphs is an unsolved puzzle. Simulations by Mac Low & Ferrara (1999) indicate that gas loss through starbursts becomes efficient only at masses  $< 10^{-6}$ , an order of magnitude less than the typical mass of a dSph. The observed upper HI limits are lower than even what is expected from evolutionary mass loss from red giants in dSphs. The lack of gas is particularly puzzling in dSphs and dEs with pronounced intermediate-age populations (Carina: 3 Gyr, Hurley-Keller et al. 1998; Leo I: 2 Gyr, Gallart et al. 1999; NGC 147, Han et al. 1997), or very recent star formation (Fornax:  $\sim 200$  Myr, Grebel & Stetson 1999). However, Carignan (1998) and Carignan (1999) found extended gas lobes around the Sculptor dSph, whose radial velocities are similar to the stellar radial velocity of this galaxy. A recent re-investigation of the Leiden-Dwingeloo survey led to the detection of similar gas concentrations with matching radial velocities outside the optical boundaries of several other dSphs and to non-detections for others (Blitz & Robishaw 2000). These authors suggest that tidal effects are the most likely agent for the displacement of the gas.

## 8. STAR FORMATION IN LOCAL GROUP DWARFS

As illustrated in earlier reviews (Grebel 1997, 1998, Mateo 1998, Grebel 1999) dwarf galaxies vary widely in their star formation histories, age distribution, and enrichment history. No two dwarfs are alike even within the same morphological type. In spite of this diversity, galaxy mass and proximity to a massive spiral appear to play a defining role in dwarf galaxy evolution.

### 8.1. MODES OF STAR FORMATION

The following modes of star formation are represented among LG dwarfs:

- Continuous star formation with a constant or varying star formation rate over a Hubble time and gradual enrichment (see also Hunter 1997). Examples for this mode include irregulars and dIrrs such as the Magellanic Clouds, which are massive enough to have a sufficiently large gas supply and to hold on to gas and metals.
- Continuous star formation with decreasing star formation rate that ceases eventually. A good example is the Fornax dSph (e.g., Grebel & Stetson 1999). This mode may be dominant in low-mass dIrrs and transition-type galaxies as well. External effects such as tidal or ram-pressure stripping may have contributed to the gradual loss of star-forming material in these galaxies. Extreme cases at the low-mass end are two of the closest and least massive Milky Way dSph com-

panions, Draco and Ursa Minor, which are dominated by ancient populations (see also Section 5). However, the abundance spread found in Draco (Shetrone et al. 1998) and the presence of a few Carbon stars indicates that the early star formation episodes must have been fairly extended.

- Distinct star formation episodes separated by Gyr-long periods of quiescence. So far only one example of this mode is known, the Carina dSph (e.g., Hurley-Keller et al. 1998). It is unclear what caused the gaps and the subsequent onset of star formation. One possible explanation is the episodic scenario proposed by Lin & Murray 1999: star formation in a dwarf heats and disperses the gas, which remains bound and eventually cools and contracts during the apogalactic passages of the dwarf around the massive parent. According to their simulations this leads to star formation separated by gaps of a few Gyr. — The distinct episodes of star formation in Carina seem to have proceeded without enrichment, a possible indication of star formation through accretion or else metal loss as described by Mac Low & Ferrara (1999).

Our present lack of knowledge of dwarf galaxy orbits makes it impossible to evaluate the impact of past close encounters and interactions on dwarf galaxy star formation histories.

### 8.2. EVOLUTION FROM dIrrs TO dSPHS

#### Star Formation Histories of MW dSph Galaxies




Figure 3. Star formation histories of Milky Way dwarf spheroidal companions. Each population box gives a schematic representation of star formation rate (SFR) as a function of age and metallicity. The fraction of intermediate-age populations tends to increase with increasing Galactocentric distance and dwarf galaxy mass. On the other hand, the distant, isolated dSph Tucana appears to have predominantly old populations.

For the Milky Way dSph companions there is a general tendency for increasing intermediate-age fractions with increasing Galactocentric distance, which appears to support ram pressure and tidal stripping (Einasto et al. 1974; Lin & Faber 1983, van den Bergh 1994) as major forces.

All morphological types of LG dwarf galaxies tend to follow global relations between absolute magnitude, mean metallicity, and central surface brightness. The more luminous a galaxy the higher its metallicity (e.g., Grebel & Guhathakurta 1999). These trends indicate the importance of dwarf galaxy mass on its evolution and ability to retain metals.

The presence of intermediate-age or even young populations in some of the more distant dSphs, the possible detection of associated gas in the surroundings of some of the dSphs, indications of substantial mass loss (Section 7.2), morphological segregation, common trends in relations between their integrated properties, and the apparent correlation between star formation histories and Galactocentric distance all seem to support the idea that low-mass dIrrs will eventually evolve into dSphs, which may be fostered by their environment. In fact the distinction between dSphs and dIrrs may be more semantical than physical.

Similarly detailed information on the M31 dSphs, which span the same range of distances from M31 as the Milky Way dSph companions from the Milky Way, would help to evaluate this scenario (Grebel & Guhathakurta 1999).

On the other hand, this cannot account for the existence of isolated, predominantly old dSphs such as Tucana. Tucana's large distance from the Milky Way and M31 (see Table 2) lets it appear unlikely that this dSph suffered major interactions in the past.

## 9. SUMMARY

Star formation histories in the LG differ from galaxy to galaxy and vary widely in length and times of their star formation episodes and enrichment history. In spite of their diversity galaxy mass and environment are important factors in determining the star formation histories. Old populations have been found in all LG galaxies studied in detail so far, though in some LG galaxies measurable star formation appears to have begun a few Gyr later than in others. Spatial variations in ages and abundances are observed in most LG galaxies. Low-mass dwarfs tend to exhibit radial population gradients with extended old populations and centrally concentrated young stars. Global modes of star formation range from continuous (variable star formation rate or gradual decrease) to episodic star formation separated by long periods of quiescence. Again, this appears to depend partially on galaxy mass and environment. Integrated properties such as mean metallicity, central surface brightness, and absolute magnitude follow common trends for all LG dwarfs. The idea that low-mass dIrrs may eventually evolve into dSphs is consistent with

the range of observed properties of transition-type dwarfs and dSphs. Major unsolved questions include the fate of the gas in dSphs, whether dSphs have been substantially affected by mass loss, unknown satellite orbits, and the problem of the “missing” LG satellites.

## ACKNOWLEDGEMENTS

This work was supported by NASA through grant HF-01108.01-98A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. I gratefully acknowledge support from the organizers through an ESA grant.

It is a pleasure to thank Robert Braun, Kem Cook, Sandy Faber, Jay Gallagher, Raja Guhathakurta, Paul Hodge, and Igor Karachentsev for valuable discussions.

## REFERENCES

- Ajhar, E.A., Grillmair, C.J., Lauer, T.R., et al. 1996, *AJ*, 111, 1110
- Aparicio, A., 1999, in *The Stellar Content of the Local Group*, IAU Symposium 193, eds. P. Whitelock & R. Cannon, Astronomical Society of the Pacific
- Armandroff, T.E., Da Costa, G.S., 1990, *AJ*, 100, 162
- Armandroff, T.E., Da Costa, G.S., Caldwell, N., Seitzer, P., 1993, *AJ*, 106, 986
- Armandroff, T.E., Davies, J.E., & Jacoby, G.H., 1998, *AJ*, 116, 2287
- Armandroff, T.E., Davies, J.E., & Jacoby, G.H., 1999, *AJ*, 118, 1220
- Bellazzini, M., Ferraro, F.R., Buonanno, R., 1999, *MNRAS*, 307, 619
- Bianchi, L., Hutchings, J.B., Massey, P., 1996, *AJ*, 111, 2303
- Blitz, L., Robishaw, T., 2000, *ApJ*, submitted (astro-ph/0001142)
- Braun, R., Burton, W.B., 1999, *A&A*, 341, 437
- Braun, R., Burton, W.B., 2000, *A&A*, 354, 853
- Buonanno, R., Corsi, C.E., Zinn, R., et al., 1998, *ApJ*, 501, L33
- Caldwell, N., 1999, *AJ*, 118, 1230
- Carignan, C., 1999, *PASA*, 16, 18
- Carignan, C., Beaulieu, S., Côté, S., et al., 1998, *AJ*, 1690
- Chandar, R., Bianchi, L., Ford, H.C., 1999, *ApJ*, 517, 668
- Cohen, J.G., Blakeslee, J.P., 1998, *AJ*, 115, 2356
- Cole, A.A., et al., 1999, *AJ*, 118, 1657
- Cook, K.H., Mario, M., Olszewski, E.W., Vogt, S.S., Stubbs, C., Diercks, A., 1999, *PASP*, 111, 306
- Côté, P., Oke, J.B., Cohen, J.G., 1999, *AJ*, 118, 1645
- Courteau, S., van den Bergh, S., 1999, *AJ*, 118, 337
- Da Costa, G.S., Armandroff, T.E., Caldwell, N., Seitzer, P., 1996, *AJ*, 112, 2576
- Da Costa, G.S., Armandroff, T.E., Caldwell, N., Seitzer, P., 2000, *AJ*, 119, 705
- Djorgovski, S.G., Gal, R.R., Odewahn, S.C., et al. 1999, in *Wide Field Surveys in Cosmology*, ed. S. Colombi et al., Editions Frontières
- Dohm-Palmer, R.C., Skillman, E.D., Saha, A., et al., 1997, *AJ*, 114, 2527
- Dolphin, A.E., 2000, *ApJ*, 531, 804
- Dressler, A. 1980, *ApJ*, 236, 351



- Dressler, A., Oemler, A., Warrick, J.C., et al. 1997, *ApJ*, 490, 577
- Feltzing, S., Gilmore, G., Wyse, R.F.G., 1999, *ApJ*, 516, L17
- Edvardsson, B., Andersen, J., Gustafsson, B., et al., 1993, *A&A*, 273, 101
- Eggen, O.J., Lynden-Bell, D., Sandage, A.R., 1962, *ApJ*, 136, 748
- Einasto, J., Saar, E., Kaasik, A., Chernin, A.D., 1974, *Nature*, 252, 111
- Eisenhauer, F., Quirrenbach, A., Zinnecker, H., Genzel, R., 1998, *ApJ*, 498, 278
- Elson, R., Tanvir, N., Gilmore, G., et al., 1999, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific
- Evans, N.W., Wilkinson, M.I., Guhathakurta, P., et al., 2000, *ApJ Lett*, *subm.*
- Gallagher, J.S., Wyse, R.F.G., 1994, *PASP*, 106, 1225
- Gallagher, J.S., Tolstoy, E., Dohm-Palmer, R.C., et al., 1998, *AJ*, 115, 1869
- Gallart, C., Aparicio, A., Vílchez, J.M., 1996, *AJ*, 112, 1950
- Gallart, C., Freedman, W.L., Aparicio, A., et al., 1999, *AJ*, 118, 2245
- Garnett, D.R., 1999, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific
- Geisler, D., Armandroff, T.E., Da Costa, G., et al., 1999, in *The Stellar Content of the Local Group*, IAU Symposium 193, eds. P. Whitelock & R. Cannon, Astronomical Society of the Pacific
- Girardi, L., Chiosi, C., Bertelli, G., Bressan, A., 1995, *A&A*, 298, 87
- González, G., Wallerstein, G., 1999, *AJ*, 117, 2286
- Grebel, E.K., 1997, *Rev. in Mod. Astron.*, 10, 29
- Grebel, E.K., 1999, in *The Stellar Content of the Local Group*, IAU Symposium 193, eds. P. Whitelock & R. Cannon, Astronomical Society of the Pacific
- Grebel, E.K., 2000, in *Massive Stellar Clusters*, Workshop of the Observatoire de Strasbourg, eds. A. Lançon & C. Boily, Astronomical Society of the Pacific
- Grebel, E.K., Brandner, W., 1999, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific
- Grebel, E.K., Guhathakurta, P., 1999, *ApJ*, 511, L101
- Grebel, E.K., Stetson, P.B., 1999, in *The Stellar Content of the Local Group*, IAU Symposium 193, eds. P. Whitelock & R. Cannon, Astronomical Society of the Pacific
- Grebel, E.K., Zaritsky, D., Harris, J., Thompson, I., 1999, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific
- Grebel, E.K., Chu, Y.-H., 2000, *AJ*, 119, 787
- Grillmair, C.J., Lauer, T.R., Worthey, G., et al., 1996, *AJ*, 112, 1975
- Grillmair, C.J., Mould, J.R., Holtzman, J.A., et al. 1998, *AJ*, 115, 144
- Guhathakurta, P., Grebel, E.K., Pittroff, L.C., et al., 2000, *AJ*, in prep.
- Gunn, J.E., Weinberg, D.H., 1995, in *Wide Field Spectroscopy and the Distant Universe*, ed. S. Maddox & A. Aragón-Salamanca, World Scientific
- Ferguson, H.C., Binggeli, B., 1994, *A&ARv*, 6, 67
- Figer, D.F., Kim, S.S., Morris, M., et al., 1999, *ApJ*, 525, 750
- Freedman, W.L., Madore, B.F., 1990, *ApJ*, 365, 186
- Freedman, W.L., Wilson, C.D., Madore, B.F., 1991, *ApJ*, 372, 455
- Han, M., Hoessel, J.G., Gallagher, J.S., et al., 1997, *AJ*, 113, 1001
- Hatzidimitriou, D., Cannon, R.D., Hawkins, M.R.S., 1993, *MNRAS*, 261, 873
- Held, E.V., Saviane, I., Momany, Y., 1999, *A&A*, 345, 747
- Held, E.V., Saviane, I., Momany, Y., Carraro, G., 2000, *ApJ*, 530, L85
- Hodge, P.W., Dolphin, A.E., Smith, T.R., Mateo, M., 1999, *ApJ*, 521, 577
- Holtzman, J.A., Watson, A.M., Baum, W.A., et al., 1998, *AJ*, 115, 1946
- Holtzman, J.A., Gallagher, J.S., Cole, A.A., et al., 1999, *AJ*, 118, 2262
- Holtzman, J.A., Mould, J.R., Gallagher, J.S., 1999b, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific
- Hunter, D.A., 1997, *PASP*, 109, 937
- Hurley-Keller, D., Mateo, M., Nemec, J., 1998, *AJ*, 115, 1840
- Hurley-Keller, D., Mateo, M., Grebel, E.K., 1999, *ApJ*, 523, L25
- Ibata, R.A., Gilmore, G., Irwin, M.J., 1994, *Nature*, 370, 194
- Ibata, R.A., Irwin, M.J., Lewis, G., Stolte, A., 2000, *ApJ*, submitted (astro-ph/0004255)
- Kaluzny, J., Kubiak, M., Szymanski, M., et al. 1995, *A&AS*, 112, 407
- Karachentsev, I.D., Karachentseva, V.E., Dolphin, A.E., et al., 2000, *A&A*, submitted
- Klessen, R.S., Kroupa, P., 1998, *ApJ*, 498, 143
- Kleyna, J.T., Geller, M.J., Kenyon, S.J., et al., 1998, *AJ*, 115, 2359
- Kleyna, J.T., Geller, M.J., Kenyon, S.J., Kurtz, M., 1999, *AJ*, 117, 1275
- Klypin, A., Kravtsov, A.V., Valenzuela, O., Prada, F., 1999, *ApJ*, 522, 82
- Knapp, G.R., Kerr, F.J., Bowers, P.F., 1978, *AJ*, 83, 360
- Kormendy, J., Bender, R., 1999, *ApJ*, 522, 772
- Kunkel, W.E., 1979, *ApJ*, 228, 718
- Lauer, T.R., Faber, S.M., Ajhar, C.J., et al., 1998, *AJ*, 116, 2263
- Lavery, R.J., Seitzer, P., Walker, A.R., et al., 1996, *AAS*, 188, 0903
- Layden, A.C., Sarajedini, A., 2000, *AJ*, 119, 1760
- Lee, M.G., 1995, *AJ*, 110, 1129
- Lee, M.G., 1996, *AJ*, 112, 1438
- Lee, M.G., Freedman, W., Mateo, M., et al., 1993, *AJ*, 106, 1420
- Lee, M.G., Aparicio, A., Tikhonov, N., Byun, Y.-I., Kim, E., 1999, *AJ*, 118, 853
- Lee, M.G., Kim, S.C., 2000, *AJ*, 119, 777
- Lin, D.N.C., Faber, S.M., 1983, *ApJ*, 266, L21
- Lin, D.N.C., Murray, S.D., 1999, in *Dwarf Galaxies and Cosmology*, Editions Frontières
- Mac Low, M.-M., Ferrara, A., 1999, *ApJ*, 513, 142
- Majewski, S.R., 1999, in *Globular Clusters*, eds. C. Martínez Roger et al., Cambridge University Press
- Majewski, S.R., Ostheimer, J.C., Kunkel, W.E., et al., 1999, in *New Views of the Magellanic Clouds*, IAU Symposium 190, eds. Y.-H. Chu et al., Astronomical Society of the Pacific

- Majewski, S.R., Ostheimer, J.C., Patterson, R.J., et al., 2000, *AJ*, 199, 760
- Martínez-Delgado, D., Aparicio, A., 1998, *AJ*, 115, 1462
- Massey, P., Johnson, K.E., DeGioia-Eastwood, K., 1995a, *ApJ*, 454, 151
- Massey, P., Lang, C.C., DeGioia-Eastwood, K., Garmany, C.D., 1995b, *ApJ*, 438, 188
- Massey, P., Hunter, D., 1998, *ApJ*, 498, 180
- Mateo, M., 1998, *ARA&A*, 36, 435
- Mateo, M., Nemec, J., Irwin, M., McMahon, R., 1991, *AJ*, 101, 892
- Mateo, M., Fischer, P., Krzemiński, W. 1995, *AJ*, 110, 2166
- Mighell, K.J., 1997, *AJ*, 114, 1458
- Mighell, K.J., Rich, R.M., 1996, *AJ*, 111, 777
- Mighell, K.J., Burke, C.J., 1999, *AJ*, 118, 366
- Minniti, D., Olszewski, E.W., Liebert, J., et al., 1995, *MNRAS*, 277, 1293
- Minniti, D., Zijlstra, A.A., 1996, *ApJ*, 467, 13
- Minniti, D., Zijlstra, A.A., Alonso, M.V., 1999, *AJ*, 117, 881
- Mochejska, B.J., Kaluzny, J., Krokenberger, M., 1998, *AcA*, 48, 455
- Monkiewicz, J., Mould, J.R., Gallagher, J.S., et al., 1999, *PASP*, 111, 1392
- Montegriffo, P., Bellazzini, M., Ferraro, F.R., et al., 1998, *MNRAS*, 294, 315
- Moore, B., Ghigna, S., Governato, F., et al. 1999, *ApJ*, 524, L19
- Oh, K.S., Lin, D.N.C., Richer, H.B., 2000, *ApJ*, 531, 727
- Morrison, H.L., Mateo, M., Olszewski, E.W., et al., 2000, *AJ*, in press (astro-ph/0001492)
- Olsen, K.A.G., Hodge, P.W., Mateo, M., et al., 1998, *MNRAS*, 300, 665
- Putman, M.E., Gibson, B.K., Staveley-Smith, L., et al., 1998, *Nature*, 294, 752
- Richer, M.G., McCall, M.L., 1995, *ApJ*, 445, 642
- Rocha-Pinto, H.J., Scalo, J., Maciel, W.J., Flynn, C., 2000, *ApJ*, 531, L115
- Rolleston, W.R.J., Dufton, P.L., McErlean, N.D., Venn, K.A., 1999, *A&A*, 348, 728
- Sage, L.J., Welch, G.A., Mitchell, G.F., 1998, *ApJ*, 507, 726
- Sakai, S., Madore, B.F., Freedman, W.L., 1999, *ApJ*, 511, 671
- Sarajedini, A., Geisler, D., Harding, P., Schommer, R., 1998, *ApJ*, 508, L37
- Saviane, I., Held, E.V., Piotto, G., 1996, *A&A*, 315, 40
- Saviane, I., Held, E.V., Bertelli, G., 2000, *A&A*, 355, 56
- Searle, L., Zinn, R., 1978, *ApJ*, 225, 357
- Shetrone, M.D., Bolte, M., Stetson, P.B., 1998, *AJ*, 115, 1888
- Smecker-Hane, T.A., Stetson, P.B., Hesser, J.E., Vandenberg, D.A., 1996, in *From Stars to Galaxies*, ed. C. Leitherer et al., Astronomical Society of the Pacific
- Smecker-Hane, T.A., Mandushev, G.I., Hesser, J.E., et al., 2000, in *Spectro-Photometric Dating of Stars and Galaxies*, eds. I. Hubeny, S. Heap, & R. Cornett, Astronomical Society of the Pacific
- Smith, G.H., Holtzman, J.A., Grillmair, C.J., 2000, in prep.
- Stetson, P.B., Hesser, J.E., Smecker-Hane, T.A., 1998, *PASP*, 110, 533
- Tikhonov, N.A., 1999, in *The Stellar Content of the Local Group*, IAU Symposium 193, eds. P. Whitelock & R. Cannon, Astronomical Society of the Pacific
- Tolstoy, E., Gallagher, J.S., Cole, A.A., et al., 1998, *AJ*, 116, 1244
- Tolstoy, E., Gallagher, J.S., Greggio, L., et al., 2000, *The ESO Messenger*, 99, 16
- Trentham, N., 1998, *MNRAS*, 294, 193
- Tully, R.B., Fisher, J.R., 1987, *Nearby Galaxies Atlas*, Cambridge University Press
- Unavane, M., Wyse, R.F.G., Gilmore, G., 1996, *MNRAS*, 278, 727
- van den Bergh, S., 1994, *ApJ*, 428, 617
- van den Bergh, S., 1999a, *ApJ*, 517, L97
- van den Bergh, S., 1999b, *A&ARv*, 9, 273
- van den Bergh, S., 2000a, in *The Local Group*, Cambridge University Press
- van den Bergh, S., 2000b, *PASP*, in press (astro-ph/0001040)
- van der Marel, R.P., de Zeeuw, P.T., Rix, H.-W., Quinlan, G.D., 1997, *Nature*, 385, 610
- Weinberg, M.D., 2000, *ApJ*, 532, 922
- Westerlund, B.E., 1997, *The Magellanic Clouds*, Cambridge University Press
- Wilcots, E.M., Miller, B.W., 1998, *AJ*, 116, 2363
- Wilkinson, M.I., Evans, N.W., 1999, *MNRAS*, 310, 645
- Whiting, A.B., Hau, G.K.T., Irwin, M. 1999, *AJ*, 118, 2767
- Young, L.M., Lo, K.Y., 1996, *ApJ*, 464, L59
- Young, L.M., Lo, K.Y., 1997a, *ApJ*, 476, 127
- Young, L.M., Lo, K.Y., 1997b, *ApJ*, 490, 710
- Young, L.M., 1999, *AJ*, 117, 1758
- Young, L.M., 2000, *AJ*, 119, 188
- Zaritsky, D., Harris, J., Grebel, E.K., Thompson, I.B., 2000, *ApJ*, 534, L53

Table 2. Known and possible Local Group members within a barycentric radius of 1.2 Mpc. The galaxies are listed by subgroup (Milky Way and likely companions, M31 and likely companions, and galaxies at distances  $> 500$  kpc from one of the spirals) and sorted by increasing distance. Galaxy types follow the nomenclature of van den Bergh (1994) with addition of the letter “N” for Sgr, NGC 205, and M32 to indicate that these may be nucleated dwarfs.  $D_{\odot}$  denotes the distance from the Sun. References are listed in the following column (“Ref”: 1 = Bellazzini et al. 1999, 2 = Westerlund 1997, 3 = Mighell & Burke 1999, 4 = Grillmair et al. 1998, 5 = Kaluzny et al. 1995, 6 = Mateo et al. 1995, 7 = Mighell 1997, 8 = Saviane et al. 2000, 9 = Mighell & Rich 1996, 10 = Lee et al. 1993, 11 = Held et al. 1999, 12 = Gallart et al. 1996, 13 = Freedman & Madore 1990, 14 = Grillmair et al. 1996, 15 = Lee 1996, 16 = Da Costa et al. 2000, 17 = Armandroff et al. 1993, 18 = Han et al. 1997, 19 = Armandroff et al. 1998, 20 = Martínez-Delgado & Aparicio 1998, 21 = Freedman et al. 1991, 22 = Grebel & Guhathakurta 1999, 23 = Sakai et al. 1999, 24 = Armandroff et al. 1999, 25 = Lee 1995, 26 = Gallagher et al. 1998, 27 = Cole et al. 1999, 28 = Whiting et al. 1999, 29 = Tolstoy et al. 1998, 30 = Dolphin 2000, 31 = Saviane et al. 1996, 32 = Lee et al. 1999, 33 = Lee & Kim 2000).  $D_{M31}$  from M31, and  $D_{LG}$  from the Local Group barycenter. The barycenter was assumed to be located at  $l = 121.7^{\circ}$ ,  $b = -21.3^{\circ}$  (Courteau & van den Bergh 1999), and 462 kpc. References for the absolute  $V$  magnitudes can be found in van den Bergh (2000b). Central surface brightnesses were taken from Mateo (1998) and complemented by data from Caldwell (1999) for And V, VI, and Cas dSph, from Whiting et al. (1999) for Cetus, from Lee et al. (1999) for DDO 210, and Lauer et al. (1998) for M33 and M31. The mean metallicities were taken from a variety of sources (column “Ref”; 34 = Shetrone et al. 1998, 35 = Mateo 1998, 36 Smecker-Hane et al. 2000, 37 = Grebel & Stetson 1999, 38 = Geisler et al. 1999, 39 = Guhathakurta et al. 2000, 40 = Côté et al. 1999, 41 = Tikhonov 1999, 42 = Cook et al. 1999). These values are in part based on red giant spectra or on photometry and represent mean abundances of the dominant intermediate/old population. Due to considerable enrichment and mixed populations in some of the galaxies mean metallicities may not always a useful concept. The  $\sigma$  quoted for  $\langle [Fe/H] \rangle$  indicates the metallicity spread, not the uncertainty in the  $\langle [Fe/H] \rangle$  determination.

| Name     | Type       | l<br>[ $^{\circ}$ ] | b<br>[ $^{\circ}$ ] | $D_{\odot}$<br>[kpc] | Ref | $D_{M31}$<br>[Mpc] | $D_{LG}$<br>[Mpc] | $M_V$<br>[mag] | $\mu_V$<br>[mag arcsec $^{-2}$ ] | $\langle [Fe/H] \rangle$<br>[dex] | Ref |
|----------|------------|---------------------|---------------------|----------------------|-----|--------------------|-------------------|----------------|----------------------------------|-----------------------------------|-----|
| Galaxy   | S(B)bcI-II | 0.00                | 0.00                | $8 \pm 1$            |     | 0.77               | 0.47              | -20.9          | —                                | —                                 |     |
| Sgr      | dSph,N?    | 6.00                | -15.00              | $28 \pm 3$           | 1   | 0.78               | 0.47              | -13.8          | $25.4 \pm 0.2$                   | $-1.0 \pm 0.3$                    | 1   |
| LMC      | IrIII-IV   | 280.46              | -32.89              | $50 \pm 5$           | 2   | 0.80               | 0.49              | -18.5          | $20.7 \pm 0.1$                   | $-0.7 \pm 0.4$                    | 2   |
| SMC      | IrIV/IV-V  | 302.80              | -44.30              | $63 \pm 10$          | 2   | 0.80               | 0.49              | -17.1          | $22.1 \pm 0.1$                   | $-1.0 \pm 0.4$                    | 2   |
| UMi      | dSph       | 104.95              | 44.80               | $69 \pm 4$           | 3   | 0.75               | 0.44              | -8.9           | $25.5 \pm 0.5$                   | $-2.2 \pm 0.2$                    | 3   |
| Dra      | dSph       | 86.37               | 34.72               | $79 \pm 4$           | 4   | 0.74               | 0.44              | -8.6           | $25.3 \pm 0.5$                   | $-2.1 \pm 0.4$                    | 34  |
| Sex      | dSph       | 243.50              | 42.27               | $86 \pm 6$           | 6   | 0.83               | 0.52              | -9.5           | $26.2 \pm 0.5$                   | $-1.7 \pm 0.2$                    | 35  |
| Scl      | dSph       | 287.54              | -83.16              | $88 \pm 4$           | 5   | 0.75               | 0.45              | -9.8           | $23.7 \pm 0.4$                   | $-1.8 \pm 0.2$                    | 35  |
| Car      | dSph       | 260.11              | -22.22              | $94 \pm 5$           | 7   | 0.82               | 0.52              | -9.4           | $25.5 \pm 0.4$                   | $-2.0 \pm 0.2$                    | 36  |
| For      | dSph       | 237.29              | -65.65              | $138 \pm 8$          | 8   | 0.76               | 0.46              | -13.1          | $23.4 \pm 0.3$                   | $-1.3 \pm 0.4$                    | 37  |
| Leo II   | dSph       | 220.17              | 67.23               | $205 \pm 12$         | 9   | 0.87               | 0.57              | -10.1          | $24.0 \pm 0.3$                   | $-1.9 \pm 0.2$                    | 35  |
| Leo I    | dSph       | 225.98              | 49.11               | $270 \pm 10$         | 10  | 0.92               | 0.63              | -11.9          | $22.4 \pm 0.3$                   | $-1.5 \pm 0.3$                    | 35  |
| Phe      | dIrr/dSph  | 272.49              | -68.82              | $405 \pm 15$         | 11  | 0.85               | 0.60              | -9.8           | —                                | $-1.8 \pm 0.4$                    | 11  |
| NGC 6822 | IrIV-V     | 25.34               | -18.39              | $500 \pm 20$         | 12  | 0.91               | 0.68              | -16.0          | $21.4 \pm 0.2$                   | $-1.2 \pm 0.4$                    | 35  |
| M31      | SbI-II     | 121.18              | -21.57              | $770 \pm 40$         | 13  | 0.00               | 0.31              | -21.2          | $10.8 \pm 0.4$                   | —                                 |     |
| M32      | dE2,N      | 121.15              | -21.98              | $770 \pm 40$         | 14  | 0.00               | 0.31              | -16.5          | $11.5 \pm 0.5$                   | $-1.1 \pm 0.6$                    | 35  |
| NGC 205  | dE5p,N     | 120.72              | -21.14              | $830 \pm 35$         | 15  | 0.06               | 0.37              | -16.4          | $20.4 \pm 0.4$                   | $-0.5 \pm 0.5$                    | 38  |
| And I    | dSph       | 121.69              | -24.85              | $790 \pm 30$         | 16  | 0.05               | 0.33              | -11.8          | $24.9 \pm 0.1$                   | $-1.5 \pm 0.2$                    | 39  |
| And III  | dSph       | 119.31              | -26.25              | $760 \pm 70$         | 17  | 0.07               | 0.30              | -10.2          | $25.3 \pm 0.1$                   | $-1.5 \pm 0.2$                    | 39  |
| NGC 147  | dE5        | 119.82              | -14.25              | $755 \pm 35$         | 18  | 0.10               | 0.30              | -15.1          | $21.6 \pm 0.2$                   | $-1.1 \pm 0.4$                    | 35  |
| And V    | dSph       | 126.20              | -15.10              | $810 \pm 45$         | 19  | 0.12               | 0.36              | -9.1           | $24.8 \pm 0.2$                   | $-1.9 \pm 0.1$                    | 39  |
| And II   | dSph       | 128.87              | -29.17              | $680 \pm 25$         | 16  | 0.16               | 0.24              | -11.8          | $24.8 \pm 0.1$                   | $-1.5 \pm 0.3$                    | 40  |
| NGC 185  | dE3p       | 120.79              | -14.48              | $620 \pm 25$         | 20  | 0.17               | 0.17              | -15.6          | $20.1 \pm 0.4$                   | $-0.8 \pm 0.4$                    | 38  |
| M33      | ScII-III   | 133.61              | -31.33              | $850 \pm 40$         | 21  | 0.22               | 0.42              | -18.9          | $10.7 \pm 0.4$                   | —                                 |     |
| Cas dSph | dSph       | 109.46              | -9.94               | $760 \pm 70$         | 22  | 0.22               | 0.34              | -12.0          | $23.5 \pm 0.1$                   | $-1.6 \pm 0.2$                    | 39  |
| IC 10    | IrIV:      | 118.97              | -3.34               | $660 \pm 65$         | 23  | 0.25               | 0.26              | -16.3          | $22.1 \pm 0.4$                   | -1.3:                             | 41  |
| And VI   | dSph       | 106.01              | -36.30              | $775 \pm 35$         | 24  | 0.27               | 0.38              | -11.3          | $24.3 \pm 0.1$                   | $-1.9 \pm 0.2$                    | 39  |
| LGS 3    | dIrr/dSph  | 126.75              | -40.90              | $810 \pm 60$         | 25  | 0.28               | 0.41              | -10.5          | $24.7 \pm 0.2$                   | $-2.2 \pm 0.3$                    | 42  |
| Peg      | IrV        | 94.77               | -43.55              | $760 \pm 100$        | 26  | 0.41               | 0.44              | -12.3          | —                                | $-1.3 \pm 0.3$                    | 26  |
| IC 1613  | IrV        | 129.82              | -60.54              | $715 \pm 35$         | 27  | 0.50               | 0.47              | -15.3          | $22.8 \pm 0.3$                   | $-1.4 \pm 0.3$                    | 27  |
| Cet      | dSph       | 101.50              | -72.90              | $775 \pm 50$         | 28  | 0.68               | 0.62              | -10.1          | $25.1 \pm 0.1$                   | $-1.9 \pm 0.2$                    | 28  |
| Leo A    | IrV        | 196.90              | 52.40               | $690 \pm 60$         | 29  | 1.11               | 0.88              | -11.5          | —                                | $-1.7 \pm 0.3$                    | 29  |
| WLM      | IrIV-V     | 75.85               | -73.63              | $945 \pm 40$         | 30  | 0.84               | 0.80              | -14.4          | $20.4 \pm 0.1$                   | $-1.4 \pm 0.4$                    | 30  |
| Tuc      | dSph       | 322.01              | 47.27               | $870 \pm 60$         | 31  | 1.22               | 1.11              | -9.6           | $25.1 \pm 0.1$                   | $-1.7 \pm 0.2$                    | 21  |